

Coherent control of the waveforms of recoilless γ 3-ray photons

Vagizov F., Antonov V., Radeonychev Y., Shakhmurov R., Kocharovskaya O.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

The concepts and ideas of coherent, nonlinear and quantum optics have been extended to photon energies in the range of 10-100 kiloelectronvolts, corresponding to soft γ 3-ray radiation (the term used when the radiation is produced in nuclear transitions) or, equivalently, hard X-ray radiation (the term used when the radiation is produced by electron motion). The recent experimental achievements in this energy range include the demonstration of parametric down-conversion in the Langevin regime, electromagnetically induced transparency in a cavity, the collective Lamb shift, vacuum-assisted generation of atomic coherences and single-photon revival in nuclear absorbing multilayer structures. Also, realization of single-photon coherent storage and stimulated Raman adiabatic passage were recently proposed in this regime. More related work is discussed in a recent review. However, the number of tools for the coherent manipulation of interactions between γ 3-ray photons and nuclear ensembles remains limited. Here we suggest and implement an efficient method to control the waveforms of γ 3-ray photons coherently. In particular, we demonstrate the conversion of individual recoilless γ 3-ray photons into a coherent, ultrashort pulse train and into a double pulse. Our method is based on the resonant interaction of γ 3-ray photons with an ensemble of nuclei with a resonant transition frequency that is periodically modulated in time. The frequency modulation, which is achieved by a uniform vibration of the resonant absorber, owing to the Doppler effect, renders resonant absorption and dispersion both time dependent, allowing us to shape the waveforms of the incident γ 3-ray photons. We expect that this technique will lead to advances in the emerging fields of coherent and quantum γ 3-ray photon optics, providing a basis for the realization of γ 3-ray-photon/nuclear- ensemble interfaces and quantum interference effects at nuclear γ 3-ray transitions. © 2014 Macmillan Publishers Limited.

<http://dx.doi.org/10.1038/nature13018>
